

# *Cerastium candidissimum*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **7.7**

Moisture indicator value: **3.6**

Reaction indicator value: **7.2**

Nutrient indicator value: **3.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.88**

Disturbance frequency (herb layer): **1.83**

Disturbance severity: **0.42**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.02**

Grazing pressure: **0.32**

Soil disturbance: **0.12**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FR \(DAP\) \*Daphno-Festucetea\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S75 Eastern Mediterranean mountain hedgehog-heath](#)

Constant species of EUNIS habitats: [S75 Eastern Mediterranean mountain hedgehog-heath](#)

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub**