

# *Cerastium carinthiacum*

## Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Chamaephyte**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.2**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **2.3**

Moisture indicator value: **4.3**

Reaction indicator value: **7.3**

Nutrient indicator value: **3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.05**

Disturbance frequency (herb layer): **1.26**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.01**

Grazing pressure: **0.15**

Soil disturbance: **0.17**

## **Habitat and sociology**

EUNIS habitat

Diagnostic species of EUNIS habitats: [U26 Temperate high-mountain base-rich scree and moraine](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**