

Cerastium cerastoides

Habitus and growth type

Plant height [m]: **0.06**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **24.04**

Flower

Flowering period: **July-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.16**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **1.2**

Moisture indicator value: **7.9**

Reaction indicator value: **4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.46**

Disturbance frequency (herb layer): **1.57**

Disturbance severity: **0.24**

Disturbance severity (herb layer): **0.23**

Mowing frequency: **0.03**

Grazing pressure: **0.13**

Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**IA \(HER\) Salicetea herbaceae**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R41 Snow-bed vegetation**](#)

Constant species of EUNIS habitats: [**R41 Snow-bed vegetation**](#)

Broad habitat

Occurrence in broad habitats: **Spring, Alpine-subalpine and arctic grassland**

Distribution

Continentality: **5**

Continentality amplitude: **4**