

# ***Cerastium pumilum***

## **Habitus and growth type**

Plant height [m]: **0.11**

Life span: **Annual**

Life form: **Therophyte**

Spinescence: **not spinescent**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **22.56**

## **Flower**

Flowering period: **February-July**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.08**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Moisture indicator value: **2.3**

Reaction indicator value: **7.3**

Nutrient indicator value: **2.1**

Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **1.56**

Disturbance frequency (herb layer): **1.99**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.39**

Grazing pressure: **0.35**

Soil disturbance: **0.31**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) \*Sedo-Scleranthetea\*](#), [FJ \(TRA\) \*Stipo-Trachynietea distachyae\*](#)

### EUNIS habitat

Constant species of EUNIS habitats: [R12 Cryptogam- and annual-dominated vegetation on siliceous rock outcrops](#), [R13 Cryptogam- and annual-dominated vegetation on calcareous and ultramafic rock outcrops](#), [R1F Mediterranean annual-rich dry grassland](#), [S34 Balkan-Anatolian submontane genistoid scrub](#)

### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**