

# ***Ceratophyllum demersum***

## **Habitus and growth type**

Plant height [m]: **1.08**

Life span: **Perennial**

Life form: **Hydrophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **25.48**

## **Flower**

Flowering period: **May-September**

## **Fruit, seed and dispersal**

Seed mass [mg]: **2.74**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.2**

Moisture indicator value: **11.9**

Reaction indicator value: **7.8**

Nutrient indicator value: **7.8**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.6**

Disturbance frequency (herb layer): **1.6**

Disturbance severity: **0.27**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.05**

Grazing pressure: **0.02**

Soil disturbance: **0.02**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [\*\*NB \(POT\) Potamogetonetea\*\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*N1J Mediterranean and Black Sea moist and wet dune slack\*\*](#), [\*\*P3b Fresh-water small pleustophyte vegetation\*\*](#), [\*\*P3c Fresh-water large pleustophyte vegetation\*\*](#), [\*\*P3d Fresh-water submerged vegetation\*\*](#), [\*\*P3e Fresh-water nymphaeid vegetation\*\*](#)

Constant species of EUNIS habitats: [\*\*P3b Fresh-water small pleustophyte vegetation\*\*](#), [\*\*P3c Fresh-water large pleustophyte vegetation\*\*](#), [\*\*P3d Fresh-water submerged vegetation\*\*](#), [\*\*P3e Fresh-water nymphaeid vegetation\*\*](#)

Dominant species of EUNIS habitats: [\*\*P3d Fresh-water submerged vegetation\*\*](#), [\*\*P3e Fresh-water nymphaeid vegetation\*\*](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

## Distribution

Continentality amplitude: **8**