

Cicendia filiformis

Habitus and growth type

Plant height [m]: **0.07**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **37.67**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.01**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **7.2**

Reaction indicator value: **2.8**

Nutrient indicator value: **1.6**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.6**

Disturbance frequency (herb layer): **1.81**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.09**

Grazing pressure: **0.18**

Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OC \(ISO\) Isoëto-Nanojuncetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [Q62 Periodically exposed shore with stable, mesotrophic sediments with pioneer or ephemeral vegetation](#)

Broad habitat

Occurrence in broad habitats: **Wetland**

Distribution

Continentality: **2**

Continentality amplitude: **2**