Cistus ocymoides

Habitus and growth type

Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.7**
Temperature indicator value: **7.2**
Moisture indicator value: **3.3**
Reaction indicator value: **2.5**
Nutrient indicator value: **1.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.55**
Disturbance frequency (herb layer): **1.49**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.21**
Mowing frequency: **0**
Grazing pressure: **0.34**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CH (ULI) *Calluno-Ulicetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S62 Western acidophilous garrigue**

Broad habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Occurrence in broad habitats: **Heathland, Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz