Cistus salviifolius

Habitus and growth type

Plant height [m]: **0.54**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **8.18**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **1.1**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **8.8**
Moisture indicator value: **2.8**
Reaction indicator value: **5**
Nutrient indicator value: **2.3**
Salinity indicator value: **0.3**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.23**
Disturbance frequency (herb layer): **1.37**
Disturbance severity: **0.67**
Disturbance severity (herb layer): **0.22**
Mowing frequency: **0.02**
Grazing pressure: **0.27**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea*, FC (LAV) *Cisto-Lavanduletea stoechadis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S51 Mediterranean maquis and arborescent matorral, S62 Western acidophilous garrigue, S63 Eastern garrigue, S71 Western Mediterranean spiny heath, T3A Mediterranean lowland to submontane Pinus forest**

Constant species of EUNIS habitats: **N1G Mediterranean coniferous coastal dune forest, S51 Mediterranean maquis and arborescent matorral, S54 Thermomediterranean arid scrub, S62 Western acidophilous garrigue, S63 Eastern garrigue, S71 Western Mediterranean spiny heath, S72 Eastern Mediterranean spiny heath (phrygana), T1A Mediterranean thermophilous deciduous forest, T21 Mediterranean evergreen Quercus forest, T3A Mediterranean lowland to submontane Pinus forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz