Clinopodium menthifolium

Habitus and growth type

Plant height [m]: **0.49**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **April-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.36**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.3**
Temperature indicator value: **6.6**
Moisture indicator value: **5**
Reaction indicator value: **6.4**
Nutrient indicator value: **4.6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.08**
Disturbance frequency (herb layer): **1.33**
Disturbance severity: **0.67**
Disturbance severity (herb layer): **0.16**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.05**
Grazing pressure: **0.21**
Soil disturbance: **0.11**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **T1G Alnus cordata forest**
Constant species of EUNIS habitats: **T1G Alnus cordata forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **5**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz