Clypeola jonthlaspi

Habitus and growth type

Plant height [m]: **0.09**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Seed mass [mg]: **0.95**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **7.8**
Moisture indicator value: **2.8**
Reaction indicator value: **8**
Nutrient indicator value: **4.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.07**
Disturbance frequency (herb layer): **1.9**
Disturbance severity: **0.76**
Disturbance severity (herb layer): **0.68**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.19**
Grazing pressure: **0.33**
Soil disturbance: **0.37**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1F Mediterranean annual-rich dry grassland**

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz