Commelina communis

Habitus and growth type

Plant height [m]: **0.48**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **June-September**

Fruit, seed and dispersal

Seed mass [mg]: **8.48**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Disturbance indicator values

Disturbance frequency: **0.68**
Disturbance frequency (herb layer): **1.59**
Disturbance severity: **0.76**
Disturbance severity (herb layer): **0.59**
Mowing frequency: **0.37**
Grazing pressure: **0.19**
Soil disturbance: **0.44**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QG (EPI) *Epilobietea angustifolii***

Broad habitat

Occurrence in broad habitats: **Scrub, Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz