

Conioselinum tataricum

Habitus and growth type

Plant height [m]: **0.8**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.8**

Temperature indicator value: **3.9**

Moisture indicator value: **6.4**

Reaction indicator value: **8**

Nutrient indicator value: **7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.26**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.3**

Mowing frequency: **0.1**

Grazing pressure: **0.18**

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [BC \(BRA\) *Brachypodio pinnati-Betuletea pendulae*](#), [CM \(MOL\) *Molinio-Arrhenatheretea*](#), [CS \(MUL\) *Mulgedio-Aconitetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3H *Larix light taiga*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest**