

# *Conopodium majus*

## Habitus and growth type

Plant height [m]: **0.36**

Life span: **Perennial**

Life form: **Geophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **22.5**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **2.01**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.2**

Temperature indicator value: **5.2**

Moisture indicator value: **4.7**

Reaction indicator value: **4.2**

Nutrient indicator value: **3.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.36**

Disturbance frequency (herb layer): **1.51**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.23**

Grazing pressure: **0.24**

Soil disturbance: **0.2**

## **Habitat and sociology**

EUNIS habitat

Constant species of EUNIS habitats: [S37 Corylus avellana scrub](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Heathland, Scrub, Sparsely vegetated (incl. rock and scree)**

## **Distribution**

Continentality: **2**

Continentality amplitude: **2**