Conopodium subcarneum

Habitus and growth type

Life span: **Perennial**
Life form: **Geophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **4**
Moisture indicator value: **5**
Reaction indicator value: **4**
Nutrient indicator value: **1**

Disturbance indicator values

Disturbance frequency: **0.43**
Disturbance frequency (herb layer): **1.6**
Disturbance severity: **0.55**
Disturbance severity (herb layer): **0.33**
Mowing frequency: **0.28**
Grazing pressure: **0.35**
Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R24 Iberian summer pasture (vallicar)**, **R33 Mediterranean short moist grassland of mountains**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz