

Corispermum intermedium

Habitus and growth type

Plant height [m]: **0.32**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **21.24**

Flower

Flowering period: **June-September**

Fruit, seed and dispersal

Seed mass [mg]: **1.99**

Dispersal mode: **Anemochory**

Dispersal distance class: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **7**

Moisture indicator value: **2.5**

Reaction indicator value: **7**

Nutrient indicator value: **6**

Salinity indicator value: **1.5**

Disturbance indicator values

Disturbance frequency: **1.73**

Disturbance frequency (herb layer): **1.99**

Disturbance severity: **0.75**

Disturbance severity (herb layer): **0.75**

Mowing frequency: **0.94**

Grazing pressure: **0.14**

Soil disturbance: **0.6**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QD \(DIG\) *Digitario sanguinalis*-*Eragrostietea minoris*](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Wetland,**

Synanthropic

Distribution

Continentality: **6**

Continentality amplitude: **1**