

Cornus sanguinea

Habitus and growth type

Plant height [m]: **3.72**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Specific leaf area [mm²/mg]: **22.34**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **109.71**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Reaction indicator value: **7.5**

Nutrient indicator value: **4.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.12**

Disturbance frequency (herb layer): **1.34**
Disturbance severity: **0.67**
Disturbance severity (herb layer): **0.19**
Mowing frequency: **0.03**
Grazing pressure: **0.2**
Soil disturbance: **0.1**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [S35 Temperate and submediterranean thorn scrub](#)

Constant species of EUNIS habitats: [S35 Temperate and submediterranean thorn scrub](#), [S37 Corylus avellana scrub](#), [S38 Temperate forest clearing scrub](#), [S53 Spartium junceum scrub](#), [T11 Temperate Salix and Populus riparian forest](#), [T13 Temperate hardwood riparian forest](#), [T19 Temperate and submediterranean thermophilous deciduous forest](#), [T1A Mediterranean thermophilous deciduous forest](#), [T1E Carpinus and Quercus mesic deciduous forest](#), [T1F Ravine forest](#), [T1G Alnus cordata forest](#), [T1H Broadleaved deciduous plantation of non site-native trees](#), [T36 Temperate and submediterranean montane Pinus sylvestris-Pinus nigra forest](#), [T3C Taxus baccata forest](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **5**