# *Coronilla coronata*

## Habitus and growth type

Plant height [m]: **0.35**
Life span: **Perennial**
Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [$mm^2$/mg]: **16.6**

## Flower

Flowering period: **April-June**

## Fruit, seed and dispersal

Seed mass [mg]: **5.89**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.3**
Temperature indicator value: **5.9**
Moisture indicator value: **3.8**
Reaction indicator value: **8.3**
Nutrient indicator value: **2.3**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.24**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Disturbance frequency (herb layer): **1.44**
Disturbance severity: **0.6**
Disturbance severity (herb layer): **0.23**
Mowing frequency: **0.15**
Grazing pressure: **0.24**
Soil disturbance: **0.12**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei*, CO (ERI) *Erico-Pinetea***

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **4**
Continentality amplitude: **2**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*