

# *Corrigiola telephiifolia*

## Habitus and growth type

Plant height [m]: **0.08**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **March-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.76**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **7.7**

Moisture indicator value: **2.7**

Reaction indicator value: **3**

Nutrient indicator value: **2.7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.9**

Disturbance frequency (herb layer): **2.01**

Disturbance severity: **0.54**

Disturbance severity (herb layer): **0.52**

Mowing frequency: **0.29**

Grazing pressure: **0.26**

Soil disturbance: **0.38**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) \*Stipo-Trachynietea distachyae\*](#)

### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline)**