

# *Crassula tillaea*

## Habitus and growth type

Plant height [m]: **0.03**

Life span: **Annual**

Life form: **Therophyte**

Spinescence: **not spinescent**

## Flower

Flowering period: **January-May**

## Fruit, seed and dispersal

Seed mass [mg]: **0.01**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.2**

Reaction indicator value: **5**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.74**

Disturbance frequency (herb layer): **2.06**

Disturbance severity: **0.56**

Disturbance severity (herb layer): **0.54**

Mowing frequency: **0.36**

Grazing pressure: **0.38**

Soil disturbance: **0.33**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OC \(ISO\) Isoëto-Nanojuncetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality amplitude: **5**