

# *Crepis bellidifolia*

## Habitus and growth type

Plant height [m]: **0.09**

Life span: **Annual, Perennial**

Life form: **Hemicryptophyte, Therophyte**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.12**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7**

Moisture indicator value: **3**

Reaction indicator value: **7**

Nutrient indicator value: **3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.84**

Disturbance frequency (herb layer): **1.23**

Disturbance severity: **0.54**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.14**

Grazing pressure: **0.23**

Soil disturbance: **0.28**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [FQ \(GEN\) \*Carici-Genistetetea lobelii\*](#)

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**