# *Crepis chondrilloides*

### Habitus and growth type

Plant height [m]: **0.35**
Life span: **Perennial**
Life form: **Hemicryptophyte**

### Flower

Flowering period: **April-June**

### Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **7**
Moisture indicator value: **2**
Reaction indicator value: **8**
Nutrient indicator value: **1**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **2.44**
Disturbance frequency (herb layer): **2.44**
Disturbance severity: **0.5**
Disturbance severity (herb layer): **0.5**
Mowing frequency: **1.33**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Grazing pressure: **0.48**
Soil disturbance: **0.19**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region**

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*