

Crepis nicaeensis

Habitus and growth type

Plant height [m]: **0.57**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **7**

Moisture indicator value: **3.6**

Reaction indicator value: **6.8**

Salinity indicator value: **0.6**

Disturbance indicator values

Disturbance frequency: **0.37**

Disturbance frequency (herb layer): **1.61**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.31**

Mowing frequency: **0.25**

Grazing pressure: **0.29**

Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

Distribution

Continentality: **4**

Continentality amplitude: **2**