

# *Crepis pusilla*

## Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **11**

Moisture indicator value: **5**

Reaction indicator value: **8**

Nutrient indicator value: **6**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.4**

Disturbance frequency (herb layer): **1.94**

Disturbance severity: **0.82**

Disturbance severity (herb layer): **0.8**

Mowing frequency: **0.25**

Grazing pressure: **0.5**

Soil disturbance: **0.47**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **[OC \(ISO\) Isoëto-Nanojuncetea](#), [QE](#)**

**(POL) Polygono-Poetea annuae**

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline),  
Synanthropic**