Crepis pyrenaica

Habitus and growth type

Plant height [m]: **0.43**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **45**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **3.18**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.5**
Temperature indicator value: **2.8**
Moisture indicator value: **5.5**
Reaction indicator value: **6.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.46**
Disturbance frequency (herb layer): **1.46**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.31**
Mowing frequency: **0.21**
Grazing pressure: **0.19**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea*, CS (MUL) *Mulgedio-Aconitetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R23 Mountain hay meadow**
Constant species of EUNIS habitats: **R23 Mountain hay meadow**

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest**

Distribution

Continentality: **4**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz