Crepis sibirica

Habitus and growth type

Plant height [m]: **0.9**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.3**
Temperature indicator value: **4**
Moisture indicator value: **5.6**
Reaction indicator value: **7**
Nutrient indicator value: **6**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.17**
Disturbance frequency (herb layer): **1.39**
Disturbance severity: **0.64**
Disturbance severity (herb layer): **0.23**
Mowing frequency: **0.1**
Grazing pressure: **0.19**
Soil disturbance: **0.11**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **BB (ASA) *Asaro europaei-Abietetea sibiricae***, **CM (MOL) *Molinio-Arrhenatheretea***, **CS (MUL) *Mulgedio-Aconitetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T3H Larix light taiga**

Constant species of EUNIS habitats: **T3H Larix light taiga**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz