Crepis tectorum

Habitus and growth type

Plant height [m]: **0.44**
Life span: **Annual, Biennial or short-lived**
Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **17.2**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.27**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry, Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**
Temperature indicator value: **5.8**
Moisture indicator value: **3.9**
Reaction indicator value: **5.5**
Nutrient indicator value: **5.8**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.38**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.97**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.68**
Mowing frequency: **0.5**
Grazing pressure: **0.19**
Soil disturbance: **0.53**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QB (SIS) *Sisymbrietea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **7**
Continentality amplitude: **7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz