# *Crocus veluchensis*

### Habitus and growth type

Life span: **Perennial**
Life form: **Geophyte**

### Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Wet**

#### Ellenberg-type indicator values

Moisture indicator value: **7**

#### Disturbance indicator values

Disturbance frequency: **0.82**
Disturbance frequency (herb layer): **1.39**
Disturbance severity: **0.36**
Disturbance severity (herb layer): **0.28**
Mowing frequency: **0.1**
Grazing pressure: **0.24**
Soil disturbance: **0.09**

### Habitat and sociology

#### Syntaxon

Diagnostic species of phytosociological classes: **CT (TRI) *Juncetea trifidi***

#### EUNIS habitat

Diagnostic species of EUNIS habitats: **Q23 Relict mire of Mediterranean mountains**, **R1K Balkan and Anatolian oromediterranean dry grassland**, **T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest**
Constant species of EUNIS habitats: **T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*