Crucianella angustifolia

Habitus and growth type

Plant height [m]: **0.25**
Life span: **Annual**
Life form: **Therophyte**
Spinescence: **not spinescent**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.47**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.4**
Moisture indicator value: **3**
Reaction indicator value: **6.5**
Nutrient indicator value: **3.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.7**
Disturbance frequency (herb layer): **1.73**
Disturbance severity: **0.61**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.28**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.32**
Soil disturbance: **0.24**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FI (TUB) *Helianthemetea guttati***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S62 Western acidophilous garrigue**

Constant species of EUNIS habitats: **S62 Western acidophilous garrigue**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz