# *Crucianella imbricata*

## Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6**
Temperature indicator value: **11**
Moisture indicator value: **4**
Reaction indicator value: **8**
Nutrient indicator value: **2**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.09**
Disturbance frequency (herb layer): **1.61**
Disturbance severity: **0.71**
Disturbance severity (herb layer): **0.26**
Mowing frequency: **0.01**
Grazing pressure: **0.33**
Soil disturbance: **0.17**

## Habitat and sociology

### Broad habitat

Occurrence in broad habitats: **Forest**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*