Crypsis alopecuroides

Habitus and growth type

Plant height [m]: **0.15**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **June-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.2**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**

Ellenberg-type indicator values

Light indicator value: **8.3**
Moisture indicator value: **6.8**
Reaction indicator value: **7.5**
Salinity indicator value: **3.8**

Disturbance indicator values

Disturbance frequency: **1.7**
Disturbance frequency (herb layer): **1.85**
Disturbance severity: **0.65**
Disturbance severity (herb layer): **0.65**
Mowing frequency: **0.25**
Grazing pressure: **0.24**
Soil disturbance: **0.33**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OC (ISO) *Isoëto-Nanojuncetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Wetland, Saline vegetation, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz