

# *Crypsis schoenoides*

## Habitus and growth type

Plant height [m]: **0.14**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **June-October**

## Fruit, seed and dispersal

Seed mass [mg]: **0.23**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

### Ellenberg-type indicator values

Light indicator value: **8.3**

Moisture indicator value: **6.9**

Reaction indicator value: **7.4**

Salinity indicator value: **5.5**

### Disturbance indicator values

Disturbance frequency: **1.82**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.61**

Mowing frequency: **0.24**

Grazing pressure: **0.25**

Soil disturbance: **0.32**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DC \(CRY\) \*Crypsietea aculeatae\*](#), [OC \(ISO\) \*Isoëto-Nanojuncetea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation](#)

Constant species of EUNIS habitats: [Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation](#)

Dominant species of EUNIS habitats: [Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation](#)

### Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Wetland, Saline vegetation, Sparsely vegetated (incl. rock and scree)**