

Cynoglossum cheirifolium

Habitus and growth type

Plant height [m]: **0.25**

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **13.9**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **8.7**

Moisture indicator value: **3.5**

Reaction indicator value: **7.5**

Nutrient indicator value: **7.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.33**

Disturbance frequency (herb layer): **1.86**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.64**

Mowing frequency: **0.35**

Grazing pressure: **0.34**

Soil disturbance: **0.34**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QF \(ART\) Artemisietea vulgaris](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**