Cyperus pannonicus

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.32**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **7**
Moisture indicator value: **8**
Reaction indicator value: **8**
Nutrient indicator value: **1**
Salinity indicator value: **5.5**

Disturbance indicator values

Disturbance frequency: **1.89**
Disturbance frequency (herb layer): **1.89**
Disturbance severity: **0.66**
Disturbance severity (herb layer): **0.66**
Mowing frequency: **0.21**
Grazing pressure: **0.26**
Soil disturbance: **0.33**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DC (CRY) *Crypsietea aculeatae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation**

Dominant species of EUNIS habitats: **Q63 Periodically exposed saline shore with pioneer or ephemeral vegetation**

Broad habitat

Occurrence in broad habitats: **Wetland, Saline vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz