

# *Cytisus borysthenicus*

## Habitus and growth type

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Moisture indicator value: **5.3**

Disturbance indicator values

Disturbance frequency: **1.5**

Disturbance frequency (herb layer): **1.5**

Disturbance severity: **0.36**

Disturbance severity (herb layer): **0.36**

Mowing frequency: **0.35**

Grazing pressure: **0.13**

Soil disturbance: **0.39**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [\*\*BD \(PYR\) \*Pyrolo-Pinetea sylvestris\*\*\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*R11 Pannonian and Pontic sandy steppe\*\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest**