

Cytisus purpureus

Habitus and growth type

Plant height [m]: **0.2**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **6.69**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **7.5**

Moisture indicator value: **5**

Reaction indicator value: **7**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.23**
Disturbance frequency (herb layer): **1.27**
Disturbance severity: **0.6**
Disturbance severity (herb layer): **0.19**
Mowing frequency: **0.08**
Grazing pressure: **0.23**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CO \(ERI\) *Erico-Pinetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T36 Temperate and submediterranean montane *Pinus sylvestris*-*Pinus nigra* forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**