Daboecia cantabrica

Habitus and growth type

Plant height [m]: **0.5**
Life span: **Perennial**
Life form: **Chamaephyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.11**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **7**
Moisture indicator value: **5**
Reaction indicator value: **2.7**
Nutrient indicator value: **1.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.32**
Disturbance frequency (herb layer): **1.36**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.05**
Grazing pressure: **0.26**
Soil disturbance: **0.1**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CH (ULI) *Calluno-Ulicetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T29 Broadleaved evergreen plantation of non site-native trees**

Broad habitat

Occurrence in broad habitats: **Heathland, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz