# *Dactylorhiza saccifera*

## Habitus and growth type

Life span: **Perennial**
Life form: **Geophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

## Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **8**
Moisture indicator value: **6.5**
Reaction indicator value: **6**
Nutrient indicator value: **2**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.28**
Disturbance frequency (herb layer): **1.36**
Disturbance severity: **0.55**
Disturbance severity (herb layer): **0.24**
Mowing frequency: **0.09**
Grazing pressure: **0.19**
Soil disturbance: **0.13**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*