

Daucus involucratus

Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **11**

Moisture indicator value: **2**

Reaction indicator value: **8**

Nutrient indicator value: **5**

Salinity indicator value: **1**

Disturbance indicator values

Disturbance frequency: **0.77**

Disturbance frequency (herb layer): **1.76**

Disturbance severity: **0.78**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.18**

Grazing pressure: **0.33**

Soil disturbance: **0.29**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [S63 Eastern garrigue](#), [S72 Eastern](#)

Mediterranean spiny heath (phrygana)

Constant species of EUNIS habitats: **S63 Eastern garrigue**

Broad habitat

Occurrence in broad habitats: **Scrub**