# *Dianthus viscidus*

### Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

### Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Disturbance indicator values

Disturbance frequency: **0.77**
Disturbance frequency (herb layer): **1.86**
Disturbance severity: **0.41**
Disturbance severity (herb layer): **0.31**
Mowing frequency: **0.03**
Grazing pressure: **0.33**
Soil disturbance: **0.11**

### Habitat and sociology

#### Syntaxon

Diagnostic species of phytosociological classes: **FO (ANA) *Trifolio anatolici-Polygonetea arenastri***

#### EUNIS habitat

Diagnostic species of EUNIS habitats: **R1K Balkan and Anatolian oromediterranean dry grassland**
Constant species of EUNIS habitats: **R1K Balkan and Anatolian oromediterranean dry grassland**

#### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*