Digitaria sanguinalis

Habitus and growth type

Plant height [m]: **0.35**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **35.25**

Flower

Flowering period: **May-November**

Fruit, seed and dispersal

Seed mass [mg]: **0.56**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.6**
Moisture indicator value: **3.7**
Reaction indicator value: **5.6**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.88**
Disturbance frequency (herb layer): **2.01**
Disturbance severity: **0.81**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.81**
Mowing frequency: **0.81**
Grazing pressure: **0.16**
Soil disturbance: **0.64**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **V34 Trampled xeric grassland with annuals**
Constant species of EUNIS habitats: **V34 Trampled xeric grassland with annuals**
Dominant species of EUNIS habitats: **V34 Trampled xeric grassland with annuals**

Broad habitat

Occurrence in broad habitats: **Wetland, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz