Dipsacus fullonum

Habitus and growth type

Plant height [m]: **1.38**
Life span: **Biennial or short-lived**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **19.64**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **2.8**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.9**
Temperature indicator value: **6.2**
Reaction indicator value: **6.7**
Nutrient indicator value: **6.4**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.11**
Disturbance frequency (herb layer): **1.54**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.63**
Disturbance severity (herb layer): **0.58**
Mowing frequency: **0.7**
Grazing pressure: **0.21**
Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QF (ART) *Artemisietea vulgaris***, **QG (EPI) *Epilobietea angustifolii***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**

Distribution

Continentality: **4**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz