

# ***Distichoselinum tenuifolium***

## **Habitus and growth type**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Spinescence: **not spinescent**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **7.48**

## **Fruit, seed and dispersal**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **8**

Moisture indicator value: **4**

Reaction indicator value: **7**

Nutrient indicator value: **7**

Salinity indicator value: **0**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*FB \(ROS\) Ononido-Rosmarinetea\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Scrub**