

Dodartia orientalis

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.06**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Disturbance indicator values

Disturbance frequency: **1.91**

Disturbance frequency (herb layer): **1.91**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.57**

Mowing frequency: **0.53**

Grazing pressure: **0.27**

Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**DB \(FEP\) Festuco-Puccinellietea**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R65 Continental subsaline alluvial pasture and meadow**](#), [**S68 Semi-desert sand dune with sparse scrub**](#)

Constant species of EUNIS habitats: [**R65 Continental subsaline alluvial pasture and meadow**](#), [**S68 Semi-desert sand dune with sparse scrub**](#)

Broad habitat

Occurrence in broad habitats: **Saline vegetation, Semi-desert**