

# ***Dodartia orientalis***

## **Habitus and growth type**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.06**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

Disturbance indicator values

Disturbance frequency: **1.91**

Disturbance frequency (herb layer): **1.91**

Disturbance severity: **0.58**

Disturbance severity (herb layer): **0.57**

Mowing frequency: **0.53**

Grazing pressure: **0.27**

Soil disturbance: **0.26**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [\*\*DB \(FEP\) Festuco-Puccinellietea\*\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [\*\*R65 Continental subsaline alluvial pasture and meadow\*\*](#), [\*\*S68 Semi-desert sand dune with sparse scrub\*\*](#)

Constant species of EUNIS habitats: [\*\*R65 Continental subsaline alluvial pasture and meadow\*\*](#), [\*\*S68 Semi-desert sand dune with sparse scrub\*\*](#)

Broad habitat

Occurrence in broad habitats: **Saline vegetation, Semi-desert**