

Doronicum orientale

Habitus and growth type

Plant height [m]: **0.34**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.45**

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6**

Moisture indicator value: **5.5**

Reaction indicator value: **7**

Nutrient indicator value: **5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.02**

Disturbance frequency (herb layer): **1.3**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.12**

Mowing frequency: **0**

Grazing pressure: **0.21**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) *Carpino-Fagetetea sylvaticae*](#),
[CB \(PUB\) *Quercetea pubescentis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T33 Mediterranean mountain *Abies* forest](#),
[T38 Mediterranean montane *Cedrus* forest](#)

Constant species of EUNIS habitats: [T33 Mediterranean mountain *Abies* forest](#), [T38 Mediterranean montane *Cedrus* forest](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Forest**