Dorycnium hirsutum

Habitus and growth type

Plant height [m]: **0.38**
Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **12.91**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **4.6**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**
Temperature indicator value: **9**
Moisture indicator value: **2.3**
Reaction indicator value: **7.3**
Nutrient indicator value: **3.3**
Salinity indicator value: **0**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.32**
Disturbance frequency (herb layer): **1.31**
Disturbance severity: **0.63**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.09**
Grazing pressure: **0.25**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **N1B Mediterranean and Black Sea coastal dune scrub**, **S53 Spartium junceum scrub**
Constant species of EUNIS habitats: **N1B Mediterranean and Black Sea coastal dune scrub**, **S53 Spartium junceum scrub**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014-2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz