Draba cretica

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **7**
Moisture indicator value: **3**
Reaction indicator value: **8**
Nutrient indicator value: **5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.29**
Disturbance frequency (herb layer): **1.29**
Disturbance severity: **0.4**
Disturbance severity (herb layer): **0.34**
Mowing frequency: **0.05**
Grazing pressure: **0.27**
Soil disturbance: **0.18**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FR (DAP) *Daphno-Festucetea*, KC**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

(ASP) *Asplenietea trichomanis*

Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz