

# ***Drimia pancrati*****on**

## **Habitus and growth type**

Plant height [m]:

Life span: **Perennial**

Life form: **Geophyte**

## **Flower**

Flowering period: **August-October**

## **Fruit, seed and dispersal**

Seed mass [mg]: **2.03**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

### **Ellenberg-type indicator values**

Light indicator value: **9**

Temperature indicator value: **10**

Moisture indicator value: **1**

Reaction indicator value: **4**

Nutrient indicator value: **2**

### **Disturbance indicator values**

Disturbance frequency: **0.61**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.71**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.14**

Grazing pressure: **0.33**

Soil disturbance: **0.19**

## Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [S54 Thermomediterranean arid scrub](#)

Constant species of EUNIS habitats: [S54 Thermomediterranean arid scrub](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**