

Drimia pancrati

Habitus and growth type

Plant height [m]:

Life span: **Perennial**

Life form: **Geophyte**

Flower

Flowering period: **August-October**

Fruit, seed and dispersal

Seed mass [mg]: **2.03**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **10**

Moisture indicator value: **1**

Reaction indicator value: **4**

Nutrient indicator value: **2**

Disturbance indicator values

Disturbance frequency: **0.61**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.71**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.14**

Grazing pressure: **0.33**

Soil disturbance: **0.19**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [S54 Thermomediterranean arid scrub](#)

Constant species of EUNIS habitats: [S54 Thermomediterranean arid scrub](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**