Drymocallis rupestris

Habitus and growth type

Plant height [m]: **0.34**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **20.55**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.3**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**
Moisture indicator value: **3.2**
Reaction indicator value: **6.2**
Nutrient indicator value: **1.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.83**
Disturbance frequency (herb layer): **1.54**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.47**
Disturbance severity (herb layer): **0.38**
Mowing frequency: **0.35**
Grazing pressure: **0.26**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CJ (COR) *Koelerio-Corynephoretea canescentis*, CL (GER) *Trifolio-Geranietea sanguinei***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1J Cyrno-Sardean oromediterranean siliceous dry grassland**

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Spring, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **5**
Continentality amplitude: **3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz